

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

19



Europäisches Patentamt
European Patent Office
Office européen des brevets

11 Publication number:

0 386 683
A2

IK
BPA

12

EUROPEAN PATENT APPLICATION

21 Application number: 90104229.1

51 Int. Cl.⁵: C07D 473/04, C07D 473/06,
A61K 31/52

22 Date of filing: 05.03.90

30 Priority: 10.03.89 IT 1973089

43 Date of publication of application:
12.09.90 Bulletin 90/37

64 Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE


71 Applicant: POLI INDUSTRIA CHIMICA S.p.A.
Piazza Agrippa, 1
I-20141 Milano(IT)

72 Inventor: Poli, Stefano
Piazza Agrippa, 1
I-20141 Milano(IT)
Inventor: Del Corona, Lucio
Piazza Agrippa, 1
I-20141 Milano(IT)
Inventor: Coppi, Germano
Piazza Agrippa, 1
I-20141 Milano(IT)

74 Representative: Minoja, Fabrizio
Studio Consulenza Brevettuale Via Rossini,
8
I-20122 Milano(IT)

54 Xanthine derivatives having bronchodilating activity, a process for the preparation thereof and pharmaceutical compositions containing them.

57 Xanthine derivatives of general formula I




(I)

EP 0 386 683 A2

having valuable bronchodilating activity, a process for the preparation thereof and pharmaceutical compositions containing them.

XANTHINE DERIVATIVES HAVING BRONCHODILATING ACTIVITY, A PROCESS FOR THE PREPARATION THEREOF AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM.


The present invention relates to xanthine derivatives of general formula (I)



wherein R_1 is H or C_1 - C_6 alkyl; R_2 is C_1 - C_6 alkyl, C_2 - C_6 alkoxyalkyl, C_3 - C_6 cycloalkyl, C_4 - C_7 cycloalkyl-methyl; R_3 is H, C_1 - C_6 alkyl, a C_4 - C_6 aliphatic cyclic amino residue having one or more nitrogen atoms directly linked to the xanthine imidazole ring.


The present invention also relates to the process for the preparation of compounds of formula (I) and to pharmaceutical compositions containing them.

It has been found that changing R_1 , R_2 and R_3 groups and above all introducing the cyclopropane cyclic group



compounds having a remarkably higher bronchodilating activity than that of Theophylline can be obtained, which compounds, therefore, can advantageously be used in therapy to relieve or prevent bronchial asthma symptoms and for the treatment of bronchospasm associated with chronic bronchitis and emphysema.

The compounds of the present invention are prepared by conventional methods, e. g. according to scheme A:



by nitrosating 6-aminouracil (II) (which is in its turn prepared by condensing the respective ureas with ethyl cyanoacetate in the presence of alkali alcoholates and alkylating with alkyl halides when $R_1 \neq H$), by means of an organic or sodium nitrite in alcoholic medium, in the presence of acids, reducing the resulting 5-nitroso-6-aminouracil (III) with sodium dithionite to give 5,6-diaminouracil (IV) which is then formylated with formic acid and treated with sodium hydroxide to give compounds (I).

The bronchodilating activity of the compounds of the present invention was evaluated by means of the

following tests, in which Theophylline was used as the control drug (I, $R_1 = R_2 = \text{CH}_3$, $R_3 = \text{H}$).

The compounds of the invention proved to be active at a dose of 10 mg/kg i.p. in the test of bronchoconstriction induced by intravenous administration of histamine in the guinea pig.

Male guinea pigs weighing 350-500 g, were anesthetized with 60 mg/kg i.p. of 6% pentobarbital sodium, and artificially vented (pump by Basile: 50 cycles/min., 6.5 ml/cycle); the tracheal cannula was connected with a "bronchospasm transducer 7020" (Basile) modified method by Konzett and Roessler) and changes in respiratory volume were registered on a writing microdynamometer (Basile).

Bronchospasm was induced by 5 $\mu\text{g/kg}$ i.v. of histamine, which caused a bronchoconstriction which is 80-90% the one obtained by complete occlusion of trachea. Such administration was repeated each 10 min. for a total of 3 times. 30 Seconds after each administration, air was insufflated into the lungs for 2 seconds.

The bronchodilating activities of the tested compounds are expressed as % change in bronchospasm obtained before treatment.

The data reported by way of example in Table 1 show that the activity of the novel compounds (ex. Ia and Ic) is substantially the same as the one of the standard drug.

Table 1.

Antibronchospastic activity, after i.p. administration, in the test of bronchoconstriction induced in guinea pig by histamine i.v. (mean \pm s.e. of 3 animals)		
Compounds	Treatment	Maximum inhibition (%)
	(mg/kg/i.p.)	(time of max. effect)
Ia	10	91,8 \pm 2,2 (10')
Ic	10	97,6 \pm 0,8 (10')
Theophylline	10	94,6 \pm 1,1 (10')

also active, at a dose of 10 mg/kg i.p., in the tests of bronchoconstriction induced by acetylcholine (20 $\mu\text{g/kg}$ i.v.) and by bradykinin (20 $\mu\text{g/kg}$ i.v.).

The tests, which were similar to the above described one, showed that the novel compounds (ex. Ia) have a higher activity than that of the standard drug (see Table 2).

Table 2.

Antibronchospastic activity, after i.p. administration, in the tests of bronchoconstriction induced by i.v. administration of acetylcholine and bradykinin in the guinea pig (mean \pm s.e. of 3 animals)			
Compounds	Treatment	Maximum inhibition (%)	
	(mg/kg/i.p.)	(time of max. effect)	
		Acetylcholine spasm	Bradykinin spasm
Ia	10	92,3 \pm 1,4 (5')	85,5 \pm 5,6 (10')
Theophylline	10	48,8 \pm 14,0 (5')	37,2 \pm 1,3 (5')

Compounds of general formula (I) proved to be active also after oral administration in the test of bronchoconstriction induced by histamine in the guinea pig. Particularly, as shown in Table 3, compound Ia is markedly more effective than the standard drug as a bronchodilating agent.

Table 3.

Antibronchospastic activity, after intraduodenal administration, in the test of bronchoconstriction induced by i.v. administration of histamine in the guinea pig (mean \pm s.e. of 3 animals).		
Compounds	Treatment	Maximum inhibition (%)
	(mg/kg/i.p.)	(time of max. effect)
Ia	31,6	86,0 \pm 11,9 (30')
	100	91,7 \pm 2,4 (30')
Theophylline	31,6	46,5 \pm 14,2 (15')
	100	94,2 \pm 1,7 (15')

The compounds of the present invention have acute toxicities equal or lower than that of Theophylline, by oral administration to the mouse.

The therapeutic index of the claimed compounds is more favourable than that of the control drug, due to the higher bronchodilating activity and the lower acute toxicity of the compounds of the invention.

Therefore, the compounds of the present invention can advantageously be used in therapy to relieve or prevent bronchial asthma symptoms and for the treatment of bronchospasm associated with chronic bronchitis and emphysema.



The present invention also relates to all of the industrially applicable aspects connected to the use of compounds of formula (I) as therapeutical agents. Therefore, an essential aspect of the invention is provided by pharmaceutical compositions containing, as the active ingredient, prefixed therapeutically effective amounts of one of compounds (I), besides possible excipients conventionally used in pharmaceutical technique. Examples of said pharmaceutical compositions comprise tablets, sugar-coated pills, hard- and soft-gelatin capsules, syrups, powders for the oral administration; lyophilized vials and solutions for the parenteral administration, and suppositories for rectal administration.

The single dosage for the oral, parenteral or rectal administration can range from 10 to 500 mg of one of compounds of formula (I), preferably from 25 to 200 mg.

The following examples further illustrate the present invention without limiting the spirit and scope thereof.

EXAMPLE 1

3-Cyclopropylmethyl-xanthine



(compound Ia)

35g (0.307 mole) of cyclopropylmethylurea (prepared from 27 g of cyclopropylnitrile by reduction with sodium and alcohol) were added to a solution of 14.4 g (0.62 g. at.) of sodium in 288 ml of absolute ethanol and the mixture was heated to ebollition and added with 40.4 g (0.357 mole) of ethyl cyanoacetate in 100

ml of ethanol, during 2 hours. The reaction mixture was refluxed for 14 hours, then water was added thereto, ethanol was evaporated off under vacuum and the aqueous solution was acidified with acetic acid, cooled to 0 °C and filtered, washing with chilled water. 35 g (62.9%) of 1-cyclopropylmethyl-6-aminouracil were obtained and dissolved in a mixture of 260 ml of ethanol, 525 ml of water and 1 ml of conc. hydrochloric acid. The solution was heated to 50 °C and treated with 25 g (0.21 mole) of isoamyl nitrite. The reaction mixture was stirred for 30 min., cooled to 10 ° - 5 °C, filtered and washed with water and ethanol to obtain 33 g (81.3%) of 1-cyclopropylmethyl-5-nitroso-6-aminouracil, which was suspended in 90 ml of ammonium hydroxide and treated with 66 g of sodium hydrosulfite at 40-50 °C until decolorization. After filtration and drying, 29 g (93.6%) of 1-cyclopropylmethyl-5,6-diaminouracil were obtained. This compound was suspended in 175 ml of 85% formic acid, stirring for 2 hours at room temperature. By addition of ether and cooling to 0 °C, 1-cyclopropylmethyl-5-amino-6-formylaminouracil precipitated which was suspended in 96 ml of 2N NaOH; the solution was heated to 90 °C for 2 hours, cooled and acidified with acetic acid, to give 25.2 g (82.9%) of 3-cyclopropylmethylxanthine, m.p. 277.9 °C, after crystallization from methanol.

15

Analysis for C ₉ H ₁₀ N ₄ O ₂			
Calc.:	C% = 52.41	H% = 4.88	N% = 27.17
Found:	C% = 52.26	H% = 4.84	N% = 27.07

20




¹H-NMR (DMSO-D₂O/TMS)

0.3-0.6 (m, 4H, -CH₂ cycl.); 1.3 (m, 1H, CH); 3.7-3.9 (m, 2H, CH₂); 7.9 (s, 1H, imid.); 10.8 (s, 1H, NH im); 13.4 (b.s., 1H, NH pyrim.)

25

By similar procedures, the compounds listed in Table 4 were obtained.

TABLE 4

Compound	R ₁	R ₂	R ₃	Mp
Ib	H	-CH ₂ -CH ₂ -O-CH ₃	H	157-8 °C
Ic	H	n-C ₃ H ₇	-N  N-CH ₃	(2.HCl)
Id	n-C ₆ H ₁₃	n-C ₆ -H ₁₃	N	142-3 °C
Ie	n-C ₅ H ₁₁	n-C ₃ H ₇	H	198-200 °C
If	H		H	288-9 °C
Ig	H	-CH ₂ 	H	277-9 °C


40

45 Claims

1. Compounds of general formula (I)

50

55




(I)

wherein R_1 is H or C_1 - C_6 alkyl; R_2 is C_1 - C_6 alkyl, C_2 - C_6 alkoxyalkyl, C_3 - C_6 cycloalkyl, C_4 - C_7 cycloalkyl-methyl; R_3 is H, C_1 - C_6 alkyl, a C_4 - C_6 aliphatic cyclic amino residue having one or more nitrogen atoms.

2. A compound as claimed in claim 1, wherein $R_1 = R_3 =$ hydrogen and $R_2 =$ cyclopropylmethyl.
3. A compound as claimed in claim 1, wherein $R_1 = R_3 =$ hydrogen and $R_2 =$ methoxyethyl.
4. A compound as claimed in claim 1, wherein $R_1 =$ hydrogen, $R_2 = n$ -propyl and $R_3 =$ 4-methyl-piperazin-1-yl.
5. A compound as claimed in claim 1, wherein $R_1 = R_2 =$ n-hexyl and $R_3 =$ hydrogen.
6. A compound as claimed in claim 1, wherein $R_1 =$ n-pentyl, $R_2 =$ n-propyl and $R_3 =$ hydrogen.
7. A compound as claimed in claim 1, wherein $R_1 = R_3 =$ hydrogen and $R_2 =$ cyclopropyl.
8. A process for the preparation of the compounds as claimed in claims 1 to 7, which process comprises nitrosating 6-aminouracil (II) by means of an organic or sodium nitrite, reducing the obtained 5-nitroso-6-aminouracil (III) with sodium dithionite to give 5,6-diaminouracil (IV), which is finally formylated with formic acid and treated with sodium hydroxide to give compound (I).
9. Pharmaceutical compositions having bronchodilating activity, containing as the active ingredient one of the compounds of claims 1 to 7, in admixture with a pharmaceutically acceptable excipient.
10. Pharmaceutical compositions as claimed in claim 9, in form of tablets, gelatin capsules, powders, syrups, drops for the oral use; lyophilized vials and solutions for the parenteral use and suppositories for the rectal use.
11. The compounds as claimed in claims 1 to 7 for use as therapeutical agents.
12. The use of the compounds as claimed in claims 1 to 7 for the preparation of pharmaceutical compositions for the therapeutical treatment of bronchial asthma and bronchospasm associated with chronic bronchitis and emphysema.

Claim for the following Contracting States:ES,GR

1. A process for the preparation of compounds of general formula I



(I)

wherein R_1 is H or C_1 - C_6 alkyl; R_2 is C_1 - C_6 alkyl, C_2 - C_6 alkoxyalkyl, C_3 - C_6 cycloalkyl, C_4 - C_7 cycloalkyl-methyl; R_3 is H, C_1 - C_6 alkyl, a C_4 - C_6 aliphatic cyclic amino residue having one or more nitrogen atoms, which process comprises nitrosating 6-aminouracil (II) by means of an organic or sodium nitrite, reducing the obtained 5-nitroso-6-aminouracil (III) with sodium dithionite to give 5,6-diaminouracil (IV), which is finally formylated with formic acid and treated with sodium hydroxide to give compound (I).

IK 



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 386 683 A3**

EUROPEAN PATENT APPLICATION

Application number: **90104229.1**

Int. Cl.⁵: **C07D 473/06, C07D 473/04, A61K 31/52**

Date of filing: **05.03.90**

Priority: **10.03.89 IT 1973089**

Date of publication of application:
12.09.90 Bulletin 90/37

Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

Date of deferred publication of the search report:
15.01.92 Bulletin 92/03


Applicant: **POLI INDUSTRIA CHIMICA S.p.A.**
Piazza Agrippa, 1
I-20141 Milano(IT)

Inventor: **Poli, Stefano**
Piazza Agrippa, 1
I-20141 Milano(IT)
Inventor: **Del Corona, Lucio**
Piazza Agrippa, 1
I-20141 Milano(IT)
Inventor: **Coppi, Germano**
Piazza Agrippa, 1
I-20141 Milano(IT)

Representative: **Minoja, Fabrizio**
Studio Consulenza Brevettuale Via Rossini,
8
I-20122 Milano(IT)

Xanthine derivatives having bronchodilating activity, a process for the preparation thereof and pharmaceutical compositions containing them.

Xanthine derivatives of general formula I



(I)

having valuable bronchodilating activity, a process for the preparation thereof and pharmaceutical compositions containing them.

EP 0 386 683 A3



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 10 4229

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	EP-A-0 011 609 (DRACO) * Claims * - - - -	1-12	C 07 D 473/06 C 07 D 473/04 A 61 K 31/52
A	EP-A-0 045 094 (DRACO) * Pages 1-5 * - - - -	1-12	
A	EP-A-0 252 043 (DRACO) * Complete specification * - - - -	1-12	
A	US-A-4 338 319 (P.G. KJELLIN et al.) * Complete specification * - - - -	1-12	
A	FR-A-2 346 353 (COOPER LABORATORIES) * Pages 48-51; claims * - - - -	1-12	
A	EP-A-0 038 784 (DRACO) * Pages 24-30; claims * - - - -	1-12	
A	EP-A-0 001 735 (DRACO) * Claims * - - - -	1-12	
A	EP-A-0 010 531 (DRACO) * Claims * - - - -	1-12	TECHNICAL FIELDS SEARCHED (Int. Cl.5)
A	DE-B-1 245 969 (VEB ARZNEIMITTELWERK DRESDEN) * Column 1, lines 1-20 * - - - - -	1	C 07 D 473/00
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of search 11 November 91	Examiner LUYTEN H.
<div>CATEGORY OF CITED DOCUMENTS</div> <div>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention</div> <div>E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document</div>			